

Tylenol (Acetaminophen) Dosage Chart

**** Tylenol can be given every 4 HOURS ** DO NOT give to children under 2 MOS OF AGE ****

| Weight (lbs) | Dose (mgs) | Infant Oral Suspension (160mg/5ml) | Children's Oral Suspension (160mg/5ml) | Chewable (80 mg tab) | Jr. Strength (160 mg tab) | Adult tab 325 mg |
|--------------|------------|------------------------------------|--|----------------------|---------------------------|------------------|
| 6-11 lbs | 40 mg | 1.25 ml = ¼ tsp | --- | --- | --- | --- |
| 12-17 lbs | 80 mg | 2.5 ml = ½ tsp | 2.5 ml = ½ tsp | --- | --- | --- |
| 18-22 lbs | 120 mg | 3.75 ml = ¾ tsp | 3.75 ml = ¾ tsp | 1 ½ tabs | --- | --- |
| 23-35 lbs | 160 mg | 5 ml = 1 tsp | 5 ml = 1 tsp | 2 tabs | 1 tab | --- |
| 36-47 lbs | 240 mg | | 7.5 ml = 1½ tsp | 3 tabs | 1½ tab | --- |
| 48-59 lbs | 320 mg | | 10 ml = 2 tsp | 4 tabs | 2 tabs | 1 tab |
| 60-71 lbs | 400 mg | | 12.5 ml = 2½ tsp | 5 tabs | 2½ tabs | 1 tab |
| 72-95 lbs | 480 mg | | 15 ml = 3 tsp | 6 tabs | 3 tabs | 1½ tab |
| 96+ lbs | 640 mg | | 20 ml = 4 tsp | 8 tabs | 4 tabs | 2 tabs |

Motrin & Advil (Ibuprofen) Dosage Chart

**** Ibuprofen can be given every 6 HOURS ** DO NOT give to children under 6 MOS OF AGE ****

| Weight (lbs) | Dose (mgs) | Infant Oral Suspension (50mg/1.25ml) | Children's Oral Suspension (100mg/5ml) | Chewable (50 mg tab) | Jr. Strength (100 mg tab) | Adult tab 200 mg |
|--------------|------------|--------------------------------------|--|----------------------|---------------------------|------------------|
| 6-11 lbs | --- | --- | --- | --- | --- | --- |
| 12-17 lbs | 50 mg | 1.25 ml=1 drop | 2.5 ml = ½ tsp | --- | --- | --- |
| 18-22 lbs | 75 mg | 1.875 ml=1½ drops | 3.75 ml = ¾ tsp | 1 ½ tabs | --- | --- |
| 23-35 lbs | 100 mg | 2.5 ml=2 drops | 5 ml = 1 tsp | 2 tabs | 1 tab | --- |
| 36-47 lbs | 240 mg | | 7.5 ml = 1½ tsp | 3 tabs | 1½ tab | --- |
| 48-59 lbs | 320 mg | | 10 ml = 2 tsp | 4 tabs | 2 tabs | 1 tab |
| 60-71 lbs | 400 mg | | 12.5 ml = 2½ tsp | 5 tabs | 2½ tabs | 1 tab |
| 72-95 lbs | 480 mg | | 15 ml = 3 tsp | 6 tabs | 3 tabs | 1½ tab |
| 96+ lbs | 640 mg | | 20 ml = 4 tsp | 8 tabs | 4 tabs | 2 tabs |

Benadryl (Diphenhydramine) Dosage Chart

**** Benadryl can be given every 6 HOURS ** Consult your physician for children under 12 MOS OF AGE ****

| Weight (lbs) | Age (years) | Suspension (12.5mg/5ml) | Chewable (12.5 mg tab) | Fastmelts (12.5 mg tab) | Tab/Cap 25 mg |
|--------------|-------------|-------------------------|------------------------|-------------------------|---------------|
| 12-19 lbs | 1-1½ yrs | 2.5 ml = ½ tsp | | | |
| 20-26 lbs | 1½-2 yrs | 3.75 ml = ¾ tsp | | | |
| 27-39 lbs | 2-4 yrs | 5 ml = 1 tsp | 1 tab | 1 tab | |
| 40-52 lbs | 5-6 yrs | 7.5 ml = 1 ½ tsp | 1 ½ tab | 1 ½ tab | |
| 53-67 lbs | 7-8 yrs | 10 ml = 2 tsp | 2 tabs | 2 tabs | 1 tab/cap |
| 68-79 lbs | 9-10 yrs | 10-12.5ml = 2-2½tsp | 2-2½ tabs | 2 tabs | 1 tab/cap |
| 80-95 lbs | 11-12 yrs | 10-15 ml = 2-3 tsp | 2-3 tabs | 2-3 tabs | 1 tab/cap |
| 96+ lbs | 12+ yrs | 10-20 ml = 2-4 tsp | 2-4 tabs | 2-4 tabs | 1-2 tabs/caps |

**** CAUTION!!! Benadryl can cause significant sleepiness or an unexpected hyperactivity reaction in some children ****

**** Dosing by weight is most accurate ** Consult your physician for any questions ****